Turf and Ornamental Systemic Fungicide

For the prevention and control of turf diseases and the diseases of annual and perennial flowers, bedding plants, foliage plants, ground covers, plus deciduous and evergreen trees and shrubs.

dust or spray mist. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum or using tobacco PERSONAL PROTECTIVE EQUIPMENT (PPE)

se detergent and hot water. Keep and wash PPE separately from other laundry.

esticide gets inside. Then wash thoroughly and put on clean clothing. Users should remove clothing/PPE immediately after handling this product. As soon as possible, wash thoroughly and change into clean clothing.

Do not apply directly to water, or areas where surface water is present, or to intertidal areas below the mean high water mark. Runoff from treated

IT IS A VIOLATION OF FEDERAL LAW TO USE THIS PRODUCT IN A MANNER INCONSISTENT WITH ITS LABELING. For use only by certified applicators

Jse this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This standard contains quirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It

Oo not enter or allow entry into treated areas during the restricted entry interval (REI) of 12 hours ersonal Protective Equipment (PPE) required for early entry to treated areas that is permitted under the Worker Protection Standard and that

Chemical-resistant footwear plus socks

NON-AGRICULTURAL USE REQUIREMENTS e requirements in this box apply to uses of this product that are NOT within the scope of the Worker Protection Standard for agricultural

Pesticide Storage: Store in the original container in a dry, temperature controlled area. Do not store in a manner where cross-contamination with Pesticide Disposal: Pesticide wastes are toxic. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal Law. If

Waste representative at the nearest EPA Regional Office for guidance. Container Handling: Nonrefillable container. Do not reuse or refill this container. Completely emoty bag into application equipment. Then offer for recycling if available or dispose of empty bag in a sanitary landfill or by incineration or, if allowed by state and local authorities, by burning

COMMERCIAL TURE AND ORNAMENTA

GENERAL INFORMATION 3336 DG Lite is a broad spectrum fungicide exhibiting preventative, curative and systemic properties. It is useful on a wide variety of turf and accurately deliver this product to the application site. Do not apply by hand or with hand held rotary spreader (e.g. bellygrinder). Application higher rate under conditions of severe disease pressure. Contact your local State Extension service for recommendations for application schedules appropriate to your area. This product is not for homeowner use and can only be used by individuals/firms licensed or registered by the state to apply ornamental or turf pest control products. No claim of compatibility with other pesticides is implied. Do not apply this product through any

of action. Cleary Chemical does not recommend the use of products containing thiabendazole in combination or rotation with 3336 DG Lite. These itilize similar chemistry and mode of action and can contribute to development of disease tolerance. If, after using 3336 DG Lite as recommended the treatment is not effective, a tolerant strain of fungi may be present. Consult your local Cleary Chemical representative, your State Agricultura ungicide or disease control strategy. As long as recommended precautions are followed, 3336 DG Lite can remain useful for disease control.

For use on all fine turf applications such as commercial, residential and public (such as home lawns, parks, athletic fields, schools, and day care centers), and golf courses (greens, tees, fairways and aprons) of cool and warm season grasses such as Bentgrass, Bluegrass, Bernudagrass, Fescue, Ryegrass, St. Augustinegrass, Zoysiagrass, or their mixtures. 3336 DG Lite is not phytotoxic to any of the above mentioned grasses when used in accordance with the label. 3336 DG Lite is to be used for the prevention and control of the diseases mentioned below. 3336 DG Lite ma be applied to dry or moist leaf/soil conditions and should be activated with irrigation or natural rainfall within 24 hours for best results. If irrigated cation should be watered in by applicator with 0.1-0.25 inches of water to begin activation of product for disease control. When treating golf greens, treatment of aprons is recommended. Use the highest recommended rate under conditions of severe disease pressure. Do not graze nimals on treated turf. Do not feed clippings to livestock or poultry. Not for use on turf being grown for sale or other commercial use as sod Table 1: Maximum Individual Application Rates and Minimum Retreatment Intervals

Use Site	Maximum application rate of 3336 DG Lite	Minimum Retreatment Interval	Comments	
Residential or Public Areas	131 lb/Acre (3 lb / 1,000 sq ft)	14 days		
Golf Course Tees, Greens, Aprons	392 lb/Acre (9 lb / 1,000 sq ft)	14 days		
Golf Course Fairways –except Florida	262 lb/Acre (6 lb / 1,000 sq ft)	14 days	Excludes Florida	
Golf Course Fairways – Florida Only	131 lb/Acre (3 lb / 1,000 sq ft)	14 days	Florida Only	

Fairways – Florida Only	(3 lb / 1,000 sq ft)		
Table 2: Maximum Seasonal Applic Do not exceed the following amounts of			
Use Site	Maximum Pounds 3336 DG Lite per Acre per Season	Pounds 3336 DG Lite per 1,000 sq ft	Comments
Residential or Public Areas	523 lb	12 lb	4 Applications per year
Golf Course Tees, Greens, Aprons	1048 lb	24 lb	
Golf Course Fairways –except Florida	262 lb	6 lb	Excludes Florida
Golf Course Fairways – Florida Only	131 lb	3 lb	Florida Only

For prevention in historic areas of disease pressure, apply twice at 14 day Anthracnose, basa intervals when soil temperature reaches 60°F. For curative control, apply Colletotrichum graminicola when disease first appears and continue at 14 day intervals. Rotations with Apply in mid-July or when disease symptoms first appear and repeat at 14 Bermudagrass Decline Take-All-Patch For prevention, apply in fall before turf has stopped all growth activity. Apply Cool Season Brown Patch second application in early Spring when soil temperatures reach 55-60°F or when disease first appears. For curative action, apply when disease first appears in early spring and continue at 14 day intervals. Apply 2 treatments at 21 day intervals in late fall to early winter, with the last Coprinus psychromorbidus application made just prior to first permanent snow cover. Rotations with PCNB For prevention, use lowest rate and apply application before expected period of disease development. Apply when disease first appears and continue at 14 day intervals. Rotations with chlorothalonil, iprodione, or mancozeb (Protect™) Ascochyta Leaf Bligh Ascochyta spp. Rhizoctonia solari **Gray Leaf Spot** (Blast) Apply preventative application before expected period of disease development. Apply when disease first appears and make applications at 14 day intervals as needed. Rotations with chlorothalonil, iprodione, or mancozeb (Protect) are Apply middle-late Fall to early Winter before turf has stopped all growth activity or in spring during periods of thaw, or when disease symptoms first appear. Continue at 14 day intervals throughout anticipated disease period. Best these wastes cannot be used according to the label instructions, contact your state Pesticide or Environmental Control Agency, or the Hazardous results are achieved when first application is made before turf is completely Apply when disease first appears and make two applications at 14 day intervals. For prevention, apply in spring and fall.

These settings are presented as a guide for use. Age, condition of spreader and operator speed will cause variation in applied rates of materia

For prevention, apply 3 applications starting late April or early May using 21

or suppression, apply at 14 day intervals when disease first appears.

For prevention, apply in early June or based upon local Extension Service ommendations. Apply at 14 day intervals as needed. Rotations and/or tank

mix combinations may be used for season long disease prevention.

	Speed (MPH)	Swath (FT)		Pounds of 3336 DG Lite Applied per 1,000 Square Feet		
Spreader			Cone Set	3	6	9
Lesco Rotary	3	8	0.25	13	17	21
Cyclone Rotary	3	4	n/a	4	5 1/4	6 ½
Scotts R-8A1 Rotary	3	11	6	J ½	N	Q
The Andersons AccuPro 2000	3	11	6	K ½	0	R
The Andersons SR 2000	3	11	6	K ½	0	R
Spyker	3	5	n/a	3 5/7	4 ½	5 1/5
PrizeLAWN CRB III & Big Foot	3	5	n/a	I 1/2	M	Р
Earthway	3	7	n/a	12	15.5	18.5
Precision Broadcast	3	4	n/a	7 3/4	10 1/4	13 ½
Precision Drop	3	n/a	n/a	5 3/4	8 1/4	11 ½
Lely W (2)*	4.5	34	III-C	5	6 3/4	8 1/4
Vicon 03 Series*	4.5	30	0	20	31	40

Nursery, Greenhouse, Landscape & Interiorscape
Annual and Perennial Flowers, Bedding Plants, Foliage Plants, Ground Covers, plus Deciduous and Evergreen Trees and Shrubs
Do not use fruit or nuts or sap from treated trees as food or feed.

3336 DG Lite is a broad spectrum systemic fungicide which controls a variety of foliar, stem, and root diseases on a wide range of commercially important plants. 3336 DG Lite may be used for disease control and prevention programs on annual and perennial flowers, bedding plants, foliage plants, ground covers, and deciduous and evergreen trees and shrubs found in Nursery, Greenhouse, Interiorscape, and Landscape sites. Follow guidelines provided under general information for use of this product.

3336 DG Lite may be used as a broadcast, fly-on, surface, over the top, and media incorporation application for the diseases listed below. For surface and broadcast applications to established plants apply at the rates below. For cut flowers do not exceed the maximum application rate 0.5 b ai/acre/application. 3336 DG Lite may be applied to dry or moist soil conditions and should be activated with irrigation or natural rainfall within 24 hours for best results. If irrigated, application should be watered in by applicator with 0.1-0.25 inches of water to begin activation. For media for later use without activation until needed. Note: The "Directions For Use" of this product reflect the cumulative inputs from both historical field use and product testing programs. However, it is impossible to test this product on all species and cultivars. A preliminary trial is suggested on a small scale before a full treatment is applied to any

plant type not shown on this label but found in a similar use site with a listed disease problem. Wait 5-7 days after treatment to evaluate results. This product is not recommended for use on Swedish Ivy (Plectranthus australis), Boston Fern (Nephrolepis exhalta), and Easter Cactus (Hatiora gaertneri).

For control of soil borne diseases effecting Bulbs, Corms, Rhizomes, Roots, Crowns

or as broadcast, fly-on, surface, or over the top application after seeding Botrytis, Cylindrocladium, Fusarium, Gliocladiu landscape beds. For curative control, apply after seeding or transplanting Myrothecium, Penicillium, Ramularia, when disease first appears. Repeat every 21-28 days for adequate crop Rhizoctonia, Sclerotinia, and Thielaviopsis protection. Note: 3336 DG Lite does not control Pythium or Phytophthora

The directions for use of this product must be followed carefully. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, (1) THE GOODS DELIVERED TO YOU ARE FURNISHED "AS IS" BY MANUFACTURER OR SELLER AND (2) MANUFACTURER AND SELLER MAKE NO WARRANTIES, GUARANTEES. OR REPRESENTATIONS OF ANY KIND TO BUYER OR USER. EITHER EXPRESS OR IMPLIED. OR BY USAGE OF TRADE. STATUTOR' purpose, use, or eligibility of the product for any particular trade usage. Unintended consequences, including bu COMBINATION WITH THE GOODS. OR THE MANNER OF USE OR APPLICATION, INCLUDING WEATHER, ALL OF WHICH ARE BEYOND THE CONTRO OF MANUFACTURER OR SELLER AND ASSUMED BY BUYER OR USER. THIS WRITING CONTAINS ALL OF THE REPRESENTATIONS AND AGREEMENT: BETWEEN BLIYER MANUFACTURER AND SELLER AND NO PERSON OR AGENT OF MANUFACTURER OR SELLER HAS ANY AUTHORITY TO MAKE ANY REPRESENTATION OR WARRANTY OR AGREEMENT RELATING IN ANY WAY TO THESE GOODS

TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, MANUFACTURER OR SELLER SHALL NOT BE LIABLE FOR SPECIAL, INCIDENTAL, OR CONSEQUENTIAL DAMAGES, OR FOR DAMAGES IN THEIR NATURE OF PENALTIES RELATING TO THE GOODS SOLD, INCLUDING USE, APPLICATION, HANDLING, AND DISPOSAL TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, MANUFACTURER OR SELLER SHALL NOT BE LIABLE TO BUYER OR USER BY WAY OF INDEMNIFICATION TO BUYER OR TO CUSTOMERS OF BUYER, IF ANY, OR FOR ANY DAMAGES OR SUMS OF MONEY, CLAIMS OR DEMANDS WHATSOEVER, RESULTING FROM OR BY REASON OF, OR RISING OUT OF THE MISUSE, OR FAILURE TO FOLLOW LABEL WARNINGS OR INSTRUCTIONS FOR USE, OF THE GOODS SOLD BY MANUFACTURER OR SELLER TO BUYER. ALL SUCH RISKS SHALL BE ASSUMED BY THE BUYER, USER, OR ITS CUSTOMERS. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, BUYER'S OR USER'S EXCLUSIVE REMEDY, AND MANUFACTURER'S OR SELLER'S TOTAL LIABILITY SHALL BE FOR DAMAGES NOT EXCEEDING THE COST OF THE PRODUCT. If you do not agree with or do not accept any of the directions for use, the warranty disclaimers, or limitations of liability, do not use the product, and return it unopened to the Seller, and the purchase price will be refunded

3336 is a registered trademark of Cleary Chemicals LLC

ACCEPTED VIA NOTIFICATION LABEL NOT REVIEWED March 12, 2018 New York State Department of Environmental Conservation

Division of Materials Management Pesticide Product Registration

Classified for "RESTRICTED USE" in New York State under 6NYCRR Part 326

Doc id: 555928

336 DG LITE®

Turf and Ornamental Systemic Fungicide

For the prevention and control of turf diseases and the diseases of annual and perennial flowers, bedding plants, foliage plants, ground covers, plus deciduous and evergreen trees and shrubs.

ACTIVE INGREDIENT

Thiophanate-methyl		2.08%
OTHER INGREDIENTS		97.92%
	TOTAL	100.00%

Keep Out Of Reach Of Children CAUTION

For Chemical Spill, Leak, Fire, or Exposure, Call CHEMTREC (800) 424-9300 For Medical Emergencies Only, Call (877) 325-1840

FIRST AID

IF SWALLOWED

Call a poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by the poison control center or doctor. Do not give anything by mouth to an unconscious person.

HOTLINE NUMBER

Have the product container or label with you when calling a poison control center or doctor, or when going for treatment. You may also contact (877) 325-1840 for emergency medical treatment information.

EPA Reg. No. 1001-70 EPA Est. No. 9198-0H-1M, 9198-0H-2B, 9198-AL-001A **Underlined letter is first** letter used in run code on bag

Net Contents

Manufactured for: Cleary Chemicals LLC 11901 S. Austin Avenue Alsip, IL 60803



П

10495032



Grow a better tomorrow.