

DOC ID: 546331

ACCEPTED  
FOR REGISTRATION

QUALI-PRO

Classified for  
"RESTRICTED USE"

September 21, 2016

in New York State

New York State Department  
of Environmental Conservation  
Division of Materials Management  
Pesticide Product Registration

under 6NYCRR Part 326

**Bifenthrin I/T 7.9 F**  
*Insecticide/Termiticide*

For prevention and control of listed termites, carpenter ants and other pests of structures. To control listed pests in and around such areas as homes, commercial and industrial buildings, recreational areas, athletic fields, lawns, ornamentals, and greenhouses.

Controls pests in livestock and poultry houses.

For Use as a Termiticide: May only be used by individuals/firms licensed by the State to apply termiticide products. States may have more restrictive requirements regarding qualifications of persons using this product. Consult the pest control regulatory agency of your State prior to use of this product.

**ACTIVE INGREDIENT:**

Bifenthrin\* ..... % BY WT. 7.9%

**INERT INGREDIENTS:** ..... 92.1%**TOTAL:** ..... 100.0%

Quali-Pro Bifenthrin I/T 7.9 F contains 2/3 pound active ingredient per gallon.

\*Cis isomers 97% minimum, trans isomers 3% maximum.

Contains bifenthrin, the active ingredient used in Talstar P Professional Insecticide.

Quali-Pro Bifenthrin I/T 7.9 F is not manufactured or distributed by FMC Corp.

EPA Reg. No. 53883-365

**KEEP OUT OF REACH OF CHILDREN**  
**CAUTION**

For additional precautionary, handling, and use statements, see inside of this booklet.

Manufactured for:

**Control Solutions Inc.**

5903 Genoa Red Bluff • Pasadena, Texas 77507



EPA 20150225

## FIRST AID

<b>IF SWALLOWED:</b>	<ul style="list-style-type: none"><li>• Call a poison control center or doctor immediately for treatment advice.</li><li>• Have person sip a glass of water if able to swallow.</li><li>• Do not induce vomiting unless told to do so by the poison control center or doctor.</li><li>• Do not give anything by mouth to an unconscious person.</li></ul>
<b>IF INHALED:</b>	<ul style="list-style-type: none"><li>• Move person to fresh air.</li><li>• If person is not breathing, call 911 or an ambulance; then give artificial respiration, preferably by mouth-to-mouth if possible.</li><li>• Call a poison control center or doctor for further treatment advice.</li></ul>
<b>IF ON SKIN OR CLOTHING:</b>	<ul style="list-style-type: none"><li>• Take off contaminated clothing.</li><li>• Rinse skin immediately with plenty of water for 15-20 minutes.</li><li>• Call a poison control center or doctor for treatment advice.</li></ul>
<b>IF IN EYES:</b>	<ul style="list-style-type: none"><li>• Hold eye open and rinse slowly and gently with water 15-20 minutes.</li><li>• Remove contact lenses, if present, after the first 5 minutes; then continuing rinsing eye.</li><li>• Call a poison control center or doctor for treatment advice.</li></ul>

**Have the product container or label with you when calling a poison control center or doctor or going for treatment.** You may also contact SafetyCall® International (866) 897-8050 for emergency medical treatment information.

**Note to Physician** - This product is a pyrethroid. If large amounts have been ingested, the stomach and intestine should be evacuated. Treatment is symptomatic and supportive. Digestible fats, oils, or alcohol may increase absorption and so should be avoided.

## PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS CAUTION

Harmful if swallowed, inhaled, or absorbed through skin. Avoid contact with skin, eyes or clothing. Avoid breathing spray mist. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet. Remove contaminated clothing and wash before reuse.

All pesticide handlers (mixers, loaders, and applicators) must wear long-sleeved shirt and long pants, socks, shoes, and chemical-resistant gloves. After the product is diluted in accordance with label directions for use and/or when mixing and loading using a closed spray tank transfer system or an in-line injector system, shirt, pants, socks, shoes, and waterproof gloves are sufficient. In addition, all pesticide handlers must wear a respiratory protection device<sup>1</sup> when working in a non-ventilated space. All pesticide handlers must wear protective eye-wear when working in a non-ventilated space or when applying termiticide by rodding or sub-slab injection.

<sup>1</sup>Use one of the following: NIOSH-approved respirator with any R, P or HE filter or a NIOSH-approved respirator with an organic vapor (OV) cartridge or canister with any R, P, or HE prefilter.

When treating adjacent to an existing structure, the applicator must check the area to be treated and immediately adjacent areas of the structure for visible and accessible cracks and holes to prevent any leaks or significant exposures to persons occupying the structure. People present or residing in the structure during application must be advised to remove their pets and themselves from the structure if they see any signs of leakage. After application, the applicator is required to check for leaks. All leaks resulting in the deposition of termiticide in locations other than those prescribed on this label must be cleaned up prior to leaving the application site. Do not allow people or pets to contact contaminated areas or to reoccupy contaminated areas of the structure until the cleanup is completed.

#### **ENVIRONMENTAL HAZARDS**

This pesticide is extremely toxic to fish and aquatic invertebrates. Do not apply directly to water, to areas where surface water is present, or to intertidal areas below the mean high water mark. Drift and runoff from treated areas may be hazardous to aquatic organisms in neighboring areas. Do not contaminate water when disposing of equipment washwaters. Care should be used when spraying to avoid fish and reptile pets in/around ornamental ponds.

This product is highly toxic to bees exposed to direct treatment or residues on blooming crops or weeds. Do not apply this product or allow to drift to blooming crops if bees are visiting the treatment area.

#### **PHYSICAL AND CHEMICAL HAZARDS**

Do not apply water-based dilutions of Quali-Pro Bifenthrin I/T 7.9 F to electrical conduits, motor housings, junction boxes, switch boxes or other electrical equipment because of possible shock hazard.

## **DIRECTIONS FOR USE**

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

Do not apply a broadcast application to interior surfaces of homes.

Do not apply by air.

Do not apply in nurseries.

Do not apply this product through any kind of irrigation system.

Not for use on sod farm turf, golf course turf, or grass grown for seed.

Do not apply to pets, crops, or sources of electricity.

Firewood is not to be treated.

Use only in well-ventilated areas.

During any application to overhead areas of structure, cover surface below with plastic sheeting or similar material except for soil surfaces in crawl spaces.

Do not allow spray to contact food, foodstuffs, food-contacting surfaces or food utensils or water supplies.

Thoroughly wash dishes and food-handling utensils with soap and water if they become contaminated by application of this product.

Do not treat areas where food is exposed.

During indoor surface applications, do not allow dripping or runoff to occur.

Do not allow contact with treated surfaces by people or pets before spray has dried.

Quali-Pro Bifenthrin I/T 7.9 F will not discolor or otherwise harm surfaces that water alone will not discolor or otherwise harm.

Do not apply this product in patient rooms or in any rooms while occupied by the elderly or infirm.

Do not apply Quali-Pro Bifenthrin I/T 7.9 F in classrooms, libraries, sports venues, or other institutional facilities when they are occupied.

Quali-Pro Bifenthrin I/T 7.9 F may be applied with low-volume application equipment, including Actisol® and Micro-Injector®, for general surface, spot, crack and crevice, and deep harborage treatments.

Not for use on plants being grown for sale or other commercial use, or for commercial seed production, or for research purposes. For use on plants intended only for aesthetic purposes or climatic modifications and being grown in interior plantscapes, ornamental gardens or parks, or lawns and grounds.

Do not apply a broadcast application to interior surfaces of homes.

Application is prohibited directly into sewers or drains, or to any area like a gutter where drainage to sewers, storm drains, water bodies, or aquatic habitat can occur. Do not allow the product to enter any drain during or after application.

#### **Additional Application Restrictions for Residential Outdoor Surface and Space Sprays**

All outdoor applications must be limited to spot or crack-and-crevice treatments only, except for the following permitted uses:

1. Treatment to soil or vegetation around structures;
2. Applications to lawns, turf, and other vegetation;
3. Applications to building foundations, up to a maximum height of 3 feet.

Other than applications to building foundations, all outdoor applications to impervious surfaces such as sidewalks, driveways, patios, porches and structural surfaces (such as windows, doors, and eaves) are limited to spot and crack-and-crevice applications, only.

#### **Distributors Must Sell in Original Packages Only.**

#### **APPLICATION DIRECTIONS**

Not for use on plants being grown for sale or other commercial use, or for commercial seed production, or for research purposes. For use on plants intended only for aesthetic purposes or climatic modifications and being grown in interior plantscapes, ornamental gardens or parks, or lawn and grounds. Only use this product on plants being grown for aesthetic or climatic purposes and in interior and exterior sites including gardens, parks, lawns, and grounds, and other ornamental sites. Do not use on vegetation intended for sale or other commercial uses. Do not use on plants grown for seed production or research purposes.

Using this product in and around structures and building construction will prevent and control termite infestations.

To institute a barrier between the wood and the termites in the soil, the chemical emulsion must be effectively dispersed in the soil. It is important to remove unnecessary materials that contain cellulose and wood from around foundation walls, crawl spaces (inside of structure), and porches, and fix damaged plumbing and construction grade in order to deny termite access to moisture.

To use this product effectively, it is important that the service technician be familiar with current control practices including trenching, rodding, sub-slab injection, low-pressure spray applications, coarse fan spraying of soil surfaces, crack and crevice (void) injection, excavated soil treatment and brush and spray applications to infested or susceptible wood. Using these techniques correctly is essential to prevent or control infestations by subterranean termite species of genera *Reticulitermes*, *Zootermopsis*, *Coptotermes* and *Heterotermes*. When determining what procedures to follow, the service technician should consider certain variables. Some of the variables to consider are species biology and behavior, structure design, heating, ventilation, and air conditioning (HVAC) systems, water table, soil type and compaction, grade conditions, and the location and type of domestic water supplies and utilities.

For information concerning the most up-to-date control practices in a given region or locale, consult the local resources for structural pest control, state cooperative extensions or regulatory agencies.

### GENERAL APPLICATIONS AND RESTRICTIONS

Quali-Pro Bifenthrin I/T 7.9 F controls a wide range of listed pests on flowers, foliage plants, non-bearing fruit and nut trees, shrubs, and ornamental trees in interior and exterior landscapes including those in hotels, office buildings, shopping malls, etc., and around athletic fields, homes, institutional buildings, parks, and recreational areas. Non-bearing fruit and nut trees are those that will not produce a harvestable crop during the season of application.

Quali-Pro Bifenthrin I/T 7.9 F can be tank mixed with insect growth regulators and other pesticides. Observe all precautions and directions for use for each product. Physical compatibility may vary with different combinations of products, so prepare a small-scale (pint or quart jar) test sample for any combination not tested previously. Use proper proportions in the small-scale test to achieve the correct result.

Unless otherwise noted in the label instructions, use the procedure below for preparation of a new tank mix:

1. Add wettable powders to tank water.
2. Mix well.
3. Add liquids and flowables.
4. Mix well.
5. Add emulsifiable concentrates.
6. Mix well.

Try reversing the order of addition or increasing the amount of water if the combination is not compatible using the above order. **NOTE:** After increasing the amount of water, if the mixture is found to be compatible, it is necessary to recalibrate the sprayer for a higher volume application. Do not allow mixture to stand overnight.

#### Formula for Determining the Active Ingredient Content of the Finished Emulsion

$$\frac{(7.9)(\text{Fl. Oz. of Quali-Pro Bifenthrin I/T 7.9 F added to tank})}{(\text{Gallons of finished spray mix})(128)} = \% \text{ Active Ingredient in emulsion}$$

### SUBTERRANEAN TERMITE CONTROL – GENERAL DIRECTIONS

**Important:** Observe the following precautions to avoid contamination of public and private water supplies:

- Use anti-backflow equipment and procedures to prevent insecticide from being siphoned into water supplies.
- Do not contaminate cisterns, wells, or other water tanks by treating the soil beneath these structures.
- Do not treat soil where runoff may occur.

- Do not treat soil water saturated or frozen soil.
- Consult local and state specifications for recommended treatment practices in your area.
- If local or state specifications do not exist, consult the Federal Housing Administration (H.U.D.) guidance documents.

**Note: For the purposes of this label**, crawl spaces are defined as being inside of the structure.

**Critical Areas:** Points at which the foundation is penetrated or abuts another structure are critical areas. These include bath traps, cracks and expansion joints, utility entry points, and adjacent structures including patios, slab additions, and stairs.

### **Structures with Wells/Cisterns Inside Foundations**

Structures that contain wells or cisterns within the foundation of a structure can only be treated using the following techniques:

1. Do not treat soil while it is beneath or within the foundation or along the exterior perimeter of a structure that contains a well or cistern. The treated backfill method must be used if soil is removed and treated outside/away from the foundation. The treated backfill technique is described as follows:
  - a) Trench and remove soil to be treated onto heavy plastic sheeting or similar material or into a wheelbarrow.
  - b) Treat the soil at the rate of 4 gallons of dilute emulsion per 10 linear feet per foot of depth of the trench or 1 gallon per 1.0 cubic feet of soil. See **“Mixing Directions”** section of this label. Mix thoroughly into the soil taking care to contain the liquid and prevent runoff or spillage.
  - c) After the treated soil has absorbed the diluted emulsion, replace the soil into the trench.
2. Treat infested and/or damaged wood in place using an injection technique such as described in the **“Control of Wood-Infesting Insects in Wood”** section of this label.

### **Structures with Adjacent Wells/Cisterns and/or Other Water Bodies**

Applicators must inspect all structures with nearby water sources such as wells, cisterns, surface ponds, streams, and other bodies of water and evaluate, at a minimum, the treatment listed below prior to making an application:

1. Prior to treatment, if feasible, expose the water pipe(s) coming from the well to the structure, if the pipe(s) enter the structure within 3 feet of grade.
2. Prior to treatment, applicators are advised to take precautions to limit the risk of applying the termiticide into subsurface drains that could empty into any bodies of water. These precautions include evaluating whether application of the termiticide to the top of the footer may result in contamination of the subsurface drain. Factors such as depth to the drain system and soil type and degree of compaction should be taken into account in determining the depth of the treatment.
3. When appropriate (i.e., on the water side of the structure), the treated backfill technique (described above) can also be used to minimize off-site movement of termiticide.

Before these techniques are used close to cisterns, wells, or other bodies of water, seek advice from local, state, or federal agencies for information on treatment practices that are accepted in your area.

**Application Rate:** Use a 0.06% emulsion for subterranean termites. For other pests on the label, use specific listed rates.

**Mixing Directions:** Mix the termiticide use dilution in the following manner: Fill tank 1/4 to 1/3 full. Start pump to begin bypass agitation and place end of treating tool in tank to allow circulation through hose. Add appropriate amount of Quali-Pro Bifenthrin I/T 7.9 F. Add remaining amount of water. Let pump run and allow recirculation through the hose for 2 to 3 minutes.

Quali-Pro Bifenthrin I/T 7.9 F may also be combined into full tanks of water. If combined into full tanks of water, allow sufficient time for agitation and/or recirculation to ensure consistency of the dilution.

To prepare a 0.06% water emulsion ready to use, dilute 3 quarts of Quali-Pro Bifenthrin I/T 7.9 F with 99.25 gallons of water.

**Mixing:** Using the chart below, determine the volume of Quali-Pro Bifenthrin I/T 7.9 F and water required to produce the desired volume of finished emulsion

<b>Amount of Quali-Pro Bifenthrin I/T 7.9 F (Gallons except where noted)</b>			
<b>Emulsion Concentrate</b>	<b>Amount of Quali-Pro Bifenthrin I/T 7.9 F</b>	<b>Amount of Water</b>	<b>Desired Gallons of Finished Emulsion</b>
0.06%	1 oz.	127 oz.	1
	5 oz.	4.9	5
	10 oz.	9.9	10
	25 oz.	24.8	25
	1.5 qt.	49.6	50
	2.25 qt.	74.4	75
	3 qt.	99.25	100
	4.5 qt.	148.8	150
	6 qt.	198.5	200
0.12% *	2 oz.	126 oz.	1
	10 oz.	4.9	5
	19.5 oz.	9.8	10
	1.5 qt.	24.6	25
	3 qt.	49.2	50
	4.5 qt.	73.8	75
	6 qt.	98.5	100
	9 qt.	147.7	150
	3	197	200

Units of measure:

1 pint = 16 fluid ounces (oz.)

1 quart = 2 pints = 4 cups = 32 fluid ounces (oz.)

\* When treating for termites, use this rate only in conjunction with volume adjustments, foam applications, or underground services applications.

**Application Volume:** To provide maximum control and protection against termite infestation, apply the specified volume of the finished water emulsion and active ingredient as set forth in the directions for use section of this label. If soil will not accept the labeled application volume, the volume may be reduced provided there is a corresponding increase in concentration so that the amount of active ingredient applied to the soil remains the same.

**Note:** Large reductions of application volume reduce the ability to obtain a continuous barrier. Variance is allowed when volume and concentration are consistent with the label-directed rates and a continuous barrier can still be achieved.

The volume of the 0.12% emulsion may be reduced by 1/2 the labeled volume where desirable for pre- and post-construction applications. When the volume is reduced, the hole spacing for sub-slab injection and soil rodding may also need to be adjusted to account for lower volume dispersal of the termiticide in the soil. Consult the following **Volume Adjustment Chart** for details.

<b>VOLUME ADJUSTMENT CHART</b>		
<b>Rate (% emulsion)</b>	<b>0.06%</b>	<b>0.12%</b>
Volume allowed		
• Horizontal (gallons emulsion/10 ft <sup>2</sup> )	1.0 gallons	0.5 gallons
• Vertical (gallons emulsion/10 linear ft.)	4.0 gallons	2.0 gallons

**After Treatment:** All holes in commonly occupied areas into which material has been applied must be plugged. Plugs must be of a non-cellulose material or covered by an impervious, non-cellulose material.

#### **Foam Applications**

Quali-Pro Bifenthrin I/T 7.9 F dilution from 0.06 to 0.12% may be converted to foam with 2X - 40X expansion characteristics and used to control or prevent termite infestations.

Depending on the circumstances, foam applications may be used alone or in combination with liquid emulsion applications. Applications may be made behind veneers, piers, chimney bases, into rubble foundations, into block voids or structural voids, under slabs, stoops, porches, or to the soil in crawl spaces and other similar voids.

Foam and liquid application must be consistent with volume and active ingredient instructions in order to ensure proper application has been made. The volume and amount of active ingredient are essential to an effective treatment. At least 75% of the labeled liquid emulsion volume of product must be applied with the remaining percent delivered to appropriate areas using foam application. Refer to label and use instructions of the foam manufacturer and the foaming equipment manufacturer.

Foam applications are generally a good supplement to liquid treatments in difficult areas but may be used alone in difficult spots.

### **Application Under Slabs or to Soil in Crawl Spaces to Prevent or Control Termites**

When making applications, Quali-Pro Bifenthrin I/T 7.9 F foam can be used alone or in combination with liquid dilution. Whether applied as a dilution, foam, or some of both, the equivalent of at least 4 gallons of 0.06% dilution (4 ounces of Quali-Pro Bifenthrin I/T 7.9 F concentrate) per 10 linear feet must be applied for a vertical barrier, or at least 1 gallon of 0.06% dilution (1 ounce of Quali-Pro Bifenthrin I/T 7.9 F concentrate) per 10 square feet must be applied for a horizontal barrier. For a foam-only application, apply Quali-Pro Bifenthrin I/T 7.9 F concentrate in sufficient concentration and volume to equal 4 ounces of concentrate per 10 linear feet or 1 ounce of concentrate per 10 square feet. For example, 2 gallons of 0.12% dilution converted to foam and used to cover 10 linear feet is the equivalent of 4 gallons of 0.06% dilution per 10 linear feet.

### **Sand Barrier Installation and Treatment**

As long as termites have access to soil that has not been treated and can avoid soil that has been treated with Quali-Pro Bifenthrin I/T 7.9 F, they can build mud tubes over surfaces that have been treated. Cracks and spaces should be filled with play box or builder's sand and then treated in the same manner as soil. Follow the rates listed on the Quali-Pro Bifenthrin I/T 7.9 F label.

Re-treatment for subterranean termites can only be performed if there is clear evidence of reinfestation or disruption of the barrier due to construction, excavation, or landscaping and/or evidence of the breakdown of the termiticide barrier in the soil. These vulnerable or reinfested areas may be re-treated in accordance with application techniques described in this product's labeling. The timing and type of these re-treatments will vary depending on factors such as termite pressure, soil types, soil conditions and other factors which may reduce the effectiveness of the barrier.

Annual re-treatment of the structure is prohibited unless there is clear evidence that reinfestation or barrier disruption has occurred.

### **PRE-CONSTRUCTION SUBTERRANEAN TERMITE TREATMENT**

**Do not apply at a lower dosage and/or concentration than specified on this label for applications prior to the installation of the finished grade.** When treating foundations deeper than 4 feet, apply the termiticide as the backfill is being replaced; or if the construction contractor fails to notify the applicator to permit this, treat the foundation to a minimum depth of 4 feet after the backfill has been installed. The applicator must trench and rod into the trench or trench along the foundation walls and around pillars and other foundation elements at the rate prescribed from grade to a minimum depth of 4 feet. When the top of the footing is exposed, the applicator must treat the soil adjacent to the footing to a depth not to exceed the bottom of the footing. However, in no case should a structure be treated below the footing.

To produce effective pre-construction subterranean termite control, create vertical and/or horizontal chemically treated zones of protection using 0.06% emulsion of Quali-Pro Bifenthrin I/T 7.9 F. Follow the current edition of the Housing and Urban Development Minimum Property Standards to assure that F.H.A. termite-proofing requirements are met.

### **Horizontal Barriers**

Establish a horizontal chemical barrier wherever treated soil will be covered by a slab such as basement floors, carports, entrance platforms, footing trenches, and slab floors.

Apply 1 gallon of 0.06% dilution per 10 square feet, or use 1 fluid ounce of Quali-Pro Bifenthrin I/T 7.9 F per 10 square feet in sufficient water (no less than 1/2 gallon or more than 2 gallons) to provide a uniform treated barrier for the area being treated.

If the fill is coarse aggregate such as washed gravel, a sufficient volume of dilution must be applied to allow it to reach the soil beneath the coarse fill.

Make applications with a low-pressure spray (less than 50 psi) using a coarse spray nozzle. If foundation walls have not been installed around the treated soil and the slab will not be poured the same day as treatment, the treated soil must be covered with a waterproof barrier. Polyethylene sheeting may be used for this purpose.

### **Vertical Barriers**

Establish vertical barriers in critical areas such as along the inside of foundation walls, plumbing, bath traps, utility services and other features that will penetrate the slab.

Using a 0.06% dilution, apply 4 gallons of dilution per 10 linear feet per foot of depth or 4 fluid ounces of Quali-Pro Bifenthrin I/T 7.9 F per 10 linear feet per foot of depth from grade level to the top of the footing in sufficient water to provide a uniform treated barrier. Use not less than 2 gallons to not more than 8 gallons of water per 10 linear feet.

When trenching and rodding into the trench, or trenching, take care to ensure that the dilution reaches the top of the footing. Space the rod holes so that a continuous treated barrier is created, but not exceeding 12 inches apart. Avoid washing out the soil around the footing. Trenches should be about 6 inches wide and 6 inches deep. Mix the chemical dilution with the soil as it is being replaced in the trench. Inside vertical barriers may not be required for monolithic slabs.

When treating hollow block voids, use 2 gallons of dilution per 10 linear feet to assure that the dilution reaches the top of the footing.

Prior to each application, applicators must notify the general contractor, construction superintendent, or similar responsible party of the intended termiticide application and intended sites of application and instruct the responsible person to notify construction workers and other individuals to leave the area to be treated during application and until the termiticide is absorbed into the soil.

### **POST-CONSTRUCTION SUBTERRANEAN TERMITE TREATMENT**

For post-construction treatment, use a 0.06% dilution. Post-construction treatments shall be made by sub-slab injection, trenching and rodding into the trench, or trenching using low-pressure spray not exceeding 25 psi at the nozzle. Proper precautions should be taken to avoid soil washout around the footing.

Locate, identify, and mark wells, electrical conduits, water and sewer lines, and radiant heat pipes prior to application of Quali-Pro Bifenthrin I/T 7.9 F. Do not puncture or inject Quali-Pro Bifenthrin I/T 7.9 F into such structures.

### **Basements**

Treatment must be made by trenching and rodding into the trench or trenching at the rate of 4 gallons of dilution per 10 linear feet per foot of depth wherever the footing from grade to the bottom of the foundation is greater than 1 foot of depth. When the footer is greater than four feet below grade, the applicator may trench and rod into the trench or trench beside foundation walls at the rate designated for four feet of depth. Space rod holes to create a continuous insecticidal barrier, but in no case more than 12 inches apart. Depending on the type of soil, degree of compaction, and location of termite activity, the actual depth of treatment will differ. However, a

structure should never be treated below the footer. Sub-slab injection may be needed beside the inside of foundation walls, around conduits, piers, and pipes, beside both sides of interior footing-supported walls, and beside cracks and partition walls.

### **Crawl Spaces – Accessible**

For crawl spaces, apply vertical termiticide barriers at the rate of 4 gallons of emulsion per 10 linear feet per foot of depth from grade to the top of the footing, or if the footing is more than 4 feet below grade to a minimum depth of 4 feet. Apply by trenching and rodding into the trench, or trenching. Treat both sides of foundation and around all piers and pipes. Where physical obstructions such as concrete walkways adjacent to foundation elements prevent trenching, treatment may be made by rodding alone. When soil type and/or conditions make trenching prohibitive, rodding may be used. When the top of the footing is exposed, the applicator must treat the soil adjacent to the footing to a depth not to exceed the bottom of the footing. Read and follow the mixing and use direction section of the label if situations are encountered where the soil will not accept the full application volume.

1. Rod holes and trenches must not extend below the bottom of the footing.
2. Rod holes must be spaced so as to achieve a continuous termiticide barrier, but in no case more than 12 inches apart.
3. Trenches must be a minimum of 6 inches deep or to the bottom of the footing, whichever is less, and need not to be wider than 6 inches. When trenching in sloping (tiered) soil, the trench must be stepped to ensure adequate distribution and prevent termiticide from running off. The emulsion must be mixed with the soil as it is replaced in the trench.
4. When treating plenums or crawl spaces, turn off the air circulation system of the structure until application has been completed and all termiticide has been absorbed by the soil.

### **Crawl Spaces – Inaccessible**

For inaccessible interior areas such as areas where there is insufficient clearance between floor joists and ground surfaces to allow operator access, excavate if possible and treat according to the instruction for accessible crawl spaces. Otherwise, apply one or a combination of the following two methods:

1. To establish a horizontal barrier, apply to the soil surface 1 gallon of emulsion per 10 square feet overall using a nozzle pressure of less than 25 psi and a coarse application nozzle (e.g., Delavan Type RD Rain-drop, RD-7 or larger, or Spraying Systems Co. 8010LP TeeJet® or comparable nozzle). For an area that cannot be reached with the application wand, use one or more extension rods to make the application to the soil. Do not broadcast or powerspray with higher pressures.
2. To establish a horizontal barrier, drill through the foundation wall or through the floor above and treat the soil perimeter at a rate of 1 gallon of emulsion per 10 square feet. Drill spacing must be at intervals not to exceed 16 inches. Many states have smaller intervals, so check State regulations which may apply.

When treating plenums and crawl spaces, turn off the air circulation systems of the structure until application has been completed and all termiticide has been absorbed by the soil.

**Excavation Technique:** When treating in troublesome areas (e.g., beside fieldstone or rubble walls, beside faulty foundation walls, and around pipes and utility lines leading downward from the structure to a well or pond), apply using the following technique:

- a. Prepare a trench placing the removed soil onto heavy-weight plastic sheeting or similar water-impermeable material.

- b. Treat the soil with 4 gallons of 0.06% dilution per 10 linear feet per foot of depth of the trench. Completely mix the dilution into the soil exercising care to avoid liquid running off the sheeting.
- c. Place the treated soil back into the trench after it has absorbed the dilution.

**Attention:** Wear NIOSH-approved unvented goggles and a respirator when applying Quali-Pro Bifenthrin I/T 7.9 F in a confined area.

### **Foundations**

For applications made after the final grade is installed, the applicator must trench and rod into the trench or trench along the foundation walls and around pillars and other foundation elements at the rate prescribed from grade to the top of the footing. When the footing is more than four (4) feet below grade, the applicator must trench and rod into the trench or trench along the foundation walls at the rate prescribed to a minimum depth of four feet. The actual depth of treatment will vary depending on soil type, degree of compaction, and location of termite activity. When the top of the footing is exposed, the applicator must treat the soil adjacent to the footing to a depth not to exceed the bottom of the footing. However, in no case should a structure be treated below the footing.

### **Masonry Voids**

Drill and treat voids in multiple masonry elements of the structure extending from the structure to the soil in order to create a continuous treatment barrier in the area to be treated. Apply at a rate of 2 gallons of emulsion per 10 linear feet of footing using a nozzle pressure of less than 25 psi. When using this treatment, access holes must be drilled below the sill plate and should be as close as possible to the footing as is practical. Treatment of voids in block or rubble foundation walls must be closely examined. Applicators must inspect areas of possible runoff as a precaution against application leakage in the treated areas. Some areas may not be treatable or may require mechanical alteration prior to treatment.

All leaks resulting in the deposition of termiticide in locations other than those prescribed on this label must be cleaned up prior to leaving the application site. Do not allow people or pets to contact contaminated areas or to reoccupy the contaminated areas of the structure until the cleanup is completed.

**NOTE:** When treating behind veneer structures (walls, etc.), take proper care to not drill beyond the veneer. If concrete blocks exist behind the veneer, both can be drilled and treated simultaneously.

Quali-Pro Bifenthrin I/T 7.9 F may not be used in voids insulated with rigid foam insulation.

### **Slabs**

Create vertical barriers by trenching and rodding into the trench or trenching outside at a rate of 4 gallons of dilution per 10 linear feet per foot of depth and by sub-slab injection within the structure. Ensure an even distribution of chemical. Applications must not be made below the bottom of the footing.

Apply beside the outside of the foundation and under the slab on the inside of foundation walls where needed. Treatment of slabs may also be necessary under and beside both sides of any interior footing-supported walls, in all cracks and expansion joints, and beside one side of interior partitions. By long-rodding or grid pattern injection vertically through the slab, horizontal barriers may be created where necessary.

- a. To permit the creation of an uninterrupted insecticidal barrier, drill holes in the foundation and/or slab.
- b. For foundations that are less than or equal to 1 foot, dig a narrow trench about 6 inches wide beside the outside of the foundation walls. Do not dig beneath the bottom of the footing. As the soil is placed back into the trench, apply 4 gallons of 0.06% dilution per 10 linear feet per foot of depth to the trench and soil.

- c. Follow the rates stated above for basements for foundations that are deeper than 1 foot.
- d. A 0.06% dilution may be used to treat soil that is exposed and wood in bath traps.

**Annual re-treatment of the structure is prohibited unless there is clear evidence that reinfestation or barrier disruption has occurred.**

### FOOD HANDLING ESTABLISHMENTS

If used as a general spot, surface, or crack and crevice treatment, Quali-Pro Bifenthrin I/T 7.9 F may be applied in both food/feed and nonfood areas of food/feed-handling establishments.

Food/feed-handling establishments are any place other than private residences where exposed food/feed is held, processed, prepared or served, including areas for receiving, storing, packing (bottling, boxing, canning, wrapping), preparing, enclosed processing systems (dairies, edible oils, mills, syrups) of food and edible waste storage. Serving areas where food is exposed and the facility is in operation are also considered food areas.

Nonfood areas in which applications are allowed include entries and vestibules, floor drains (to sewers), garages, garbage rooms, lavatories, locker rooms, machine rooms, mop closets, offices, and storage (after canning or bottling).

Listed below are some of the use sites that are allowed:

- aircraft (do not use in aircraft cabins)
- apartment buildings
- bakeries
- bottling facilities
- breweries
- buses
- cafeterias
- candy plants
- canneries
- dairy product processing plants
- food manufacturing plants
- food processing plants
- food service establishments
- granaries
- grain mills
- hospitals
- hotels
- industrial buildings
- laboratories
- meat/poultry/egg processing plants
- mobile/motor homes
- nursing homes
- offices
- railcars
- restaurants
- schools
- ships
- trailers
- trucks
- vessels
- warehouses
- wineries

**General Surface Application:** Do not use this application method in food/feed-handling establishments when the facility is in operation or foods/feeds are exposed. During treatment, remove or cover all food-processing and/or handling equipment and do not apply directly to food products. All equipment, benches, shelving and other surfaces in food-processing plants, bakeries, cafeterias and other facilities which food will contact must be washed after treatment. Clean food-handling equipment or processing equipment and rinse completely with fresh clean water.

**Spot, Crack and Crevice Application:** These types of treatments can be done when the facility is operating, but food should be covered or removed from the treatment area. Do not apply directly to food.

**Foam Applications:** Converting Quali-Pro Bifenthrin I/T 7.9 F to foam will allow it to be used to treat structural voids. To produce a 0.02% to 0.06% foam concentration, dilute 0.33 to 1.0 fl. oz. of Quali-Pro Bifenthrin I/T 7.9 F per gallon of water and add the manufacturer's recommended amount of foaming agent. Before application, make sure that the foaming agent is compatible with Quali-Pro Bifenthrin I/T 7.9 F.

### INDOOR USES

In the home, all food-processing surfaces and utensils should be covered during treatment or thoroughly washed before reuse. Exposed food should be covered or removed. Do not permit humans or pets to contact treated surfaces until the spray has dried.

During any overhead applications to overhead interior areas of structures, cover surfaces below with plastic sheeting or similar materials.

Wear protective clothing, unvented goggles, gloves and respirator when applying to overhead areas or in poorly ventilated areas. Avoid touching sprayed surfaces until spray has completely dried.

Quali-Pro Bifenthrin I/T 7.9 F may be used to control ants, grocer beetles, boxelder bugs, carpet beetles, centipedes, clothes moths, cockroaches, crickets, earwigs, firebrats, flies, gnats, midges, millipedes, pillbugs, scorpions, silverfish, sowbugs, spiders, ticks, and wasps.

In structures and buildings and on modes of transportation, use a 0.02% to 0.06% suspension (0.33 to 1 fl. oz. per gallon of water) using a crack and crevice, pinstream, spot, coarse, low-pressure spray (25 psi or less) or with a paint brush.

**Indoor Treatments:** Apply where pests hide. These areas include but are not limited to baseboards, corners, storage areas, closets, around water pipes, doors and windows, attics and eaves, behind and under refrigerators, cabinets, sinks, furnaces, stoves, the underside of shelves, and drawers. Treat with a low-pressure, coarse, crack and crevice or spot spray. Pay close attention to cracks and crevices. Not for use as a space spray.

**Mixing Directions:** See mixing directions in “**Pest Control on Outside Surfaces and Around Buildings**” section.

To make a dilution for brush or spray treatments:

- Dilute Quali-Pro Bifenthrin I/T 7.9 F with water.
- Fill sprayer with the required amount of water.
- Add Quali-Pro Bifenthrin I/T 7.9 F.
- To ensure proper mixing, close sprayer and shake before use.
- Only mix the amount of solution that is necessary for treatment.

In order to achieve and/or maintain control in times of high pest pressure, re-treatment may be needed.

Repeat application should only take place if there are signs of renewed insect activity and must not exceed one application per 7 days.

**Ants:** Apply to any ant trails, around doors and windows, and other places that ants frequent.

**Bees and Wasps:** Apply to nests in late evening when these pests are at rest. Spray nests, entrances to nests and surrounding areas thoroughly. Contact as many insects as possible. Re-treat if signs of renewed activity exist.

**Boxelder Bugs, Centipedes, Earwigs, Beetles, Millipedes, Pillbugs, and Sowbugs:** Treat near doors and windows, storage areas, baseboards and other sites where these pests may be found.

**Cockroaches, Crickets, Firebrats, Scorpions, Silverfish, Spiders, and Ticks:** Use a coarse low-pressure, crack and crevice or spot spray paying close attention to cracks and crevices. Treat where pests hide. These areas include but are not limited to attics and eaves, baseboards, closets, corners, storage areas, around water pipes, doors and windows, behind and under cabinets, furnaces, refrigerators, sinks, stoves, and the underside of shelves and drawers.

### **LIVESTOCK AND POULTRY BUILDINGS AND KENNELS**

Quali-Pro Bifenthrin I/T 7.9 F may be used as a surface spray (including directed spray) and crack and crevice application for control of flies, fleas, beetles, bed bugs, mites and ticks. Usual cleaning of the structure must be completed along with the Quali-Pro Bifenthrin I/T 7.9 F application for effective control.

For the areas of the structure that are not in use, make application to floors, walls, pipes, windows, and doors where pests may be present at a rate of 0.33 to 1 oz per 1000 sq ft. Cover feed and water units prior to application to prevent contamination. Outside applications to sides and foundations are also helpful against interior infestation.

For areas of the structure that are in use, apply inside to cracks and crevices only at a rate of 0.33 to 1 oz per 1000 sq. feet. Outside applications to sides and foundation is also helpful against interior infestation. Do not apply directly to animals.

Quali-Pro Bifenthrin I/T 7.9 F will control adult flies by spraying in and around the animal structure. The application should be made to any where flies will sit including windows, inside and outside walls, fences, vegetation, and on manure in areas that cannot be cleaned.

Applications in poultry houses should be made at a rate of 0.33 to 1 fl oz per 1000 sq feet to floors, walls, poles, and cages (if used), cracks and crevices. In order to prevent unobtrusive pests, maintain a year-round treatment program. Applications should be made after each batch or cleaning, but not more than once every 8 weeks. Outside applications to sides (2-3 feet high), foundation, and ground (6-10 feet out) is also helpful against interior infestation.

Quali-Pro Bifenthrin I/T 7.9 F should be applied at a rate of 0.33 to 1 fl oz per 1000 sq feet for control of beetles in structures containing birds (including broilers for breeding) grown on refuse matter. Make application after animals are removed and during harrowing. When structure is cleaned out, apply to bare dirt or cement, sides, and exterior border, and treat new litter that is put down. Reapply after each batch of animals.

Use Quali-Pro Bifenthrin I/T 7.9 F at a rate of 0.33 to 1 fl oz per 1000 sq ft to treat the outside edge of the manure when used in caged-layer houses to control beetles. Do not apply directly to accrued waste. An application to sides, ceiling, poles, and exterior part of structure should be made. Reapply between each batch of animals.

### **RESTRICTIONS**

- Do not apply to milk rooms.
- Do not apply as a general surface spray when animals are present in structure. When animals are present, only cracks and crevices may be treated.
- Applications must be dry before allowing animals back into structure.
- Spray must be dry before applying a disinfectant.
- Cover feed and water units when making application so not to contaminate.

## LAWNS

Use Quali-Pro Bifenthrin I/T 7.9 F as a broadcast treatment. To accomplish uniform control when applying to dense grass foliage, use volumes of up to 10 gallons per 1000 square feet.

To ensure control of sub-surface pests including mole crickets using low-volume treatments, (i.e., less than 2 gallons per 1000 square feet), immediately follow the treatment with irrigation of the treated area with at least 0.25 inches of water.

### Lawn Application Rates

Under typical conditions, the application rates shown in the table below will provide control of the listed pests. Quali-Pro Bifenthrin I/T 7.9 F may, however, be applied at up to 1 fl. oz. per 1000 square feet at the discretion of the applicator. Maximum residual control requires the higher treatment rates.

<b>Pest</b>	<b>Application Rate Quali-Pro Bifenthrin I/T 7.9 F</b>
Armyworms <sup>1</sup> Cutworms <sup>1</sup> Sod Webworm <sup>1</sup>	0.18 - 0.25 fluid oz. per 1000 sq. ft.
Annual Bluegrass Weevil (Hyperodes)(Adult) <sup>2</sup> Banks Grass Mite <sup>6</sup> Billbugs (Adult) <sup>3</sup> Black Turfgrass Ataenius Adult <sup>4</sup> Centipedes Chinch Bugs <sup>5</sup> Crickets Earwigs Fleas (Adult) Grasshoppers Leafhoppers Mealybugs Millipedes Mites <sup>6</sup> Pillbugs Sowbugs	0.25 - 0.5 fluid oz. per 1000 sq. ft.
Ants Fleas (Larvae) <sup>7</sup> Imported Fire Ants <sup>8</sup> Japanese Beetle (Adult) Mole Cricket (Adult) <sup>9</sup> Mole Cricket (Nymph) <sup>10</sup> Ticks <sup>11</sup>	0.5 - 1.0 fluid oz. per 1000 sq. ft.

In New York State, this product may not be applied to any grass or turf area within 100 feet of a water body (lake, pond, river, stream, wetland, or drainage ditch).

In New York State, do make a single repeat application of Quali-Pro Bifenthrin I/T 7.9 F if there are signs of renewed insect activity, but not sooner than two weeks after the first application.

#### Notes

- <sup>1</sup> **Armyworms, Cutworms, and Sod Webworms:** Postpone irrigation or mowing for 24 hours after application to obtain the best possible control. Higher treatment rates (up to 1 fluid oz. per 1000 square feet) may be necessary if high pest pressure exists and if the grass is maintained taller than 1 inch.
- <sup>2</sup> **Annual Bluegrass Weevil (*Hyperodes*) adults:** Treatment of this species should be timed as they travel into grass areas and away from their overwintering sites. Travel usually begins when *Forsythia* is in full bloom and ends when *Cornus florida* (flowering dogwood) is in full bloom. For additional detailed information regarding treatment timing, check with your State Cooperative Extension Service.
- <sup>3</sup> **Billbug adults:** Treatment of adult billbugs should be made when they are first noticed in April and May. To optimize treatment timing, degree day models have been developed. For detailed information particular to your region, check with your State Cooperative Extension Service. Spring treatments for billbug adults will also offer control of over-wintered chinch bugs in temperate climates.
- <sup>4</sup> **Black Turfgrass *Ataenius* adults:** In order to control the 1st and 2nd generation of black turfgrass *ataenius* adults respectively, treatments should take place in May and July. Time the May treatment to match with the full bloom stage of *Vanhoutte spiraea* (*Spiraea vanhoutte*) and horse chestnut (*Aesculus hippocastanum*). Time the July treatment to match with the blooming Rose of Sharon (*Hibiscus syriacus*).
- <sup>5</sup> **Chinch Bugs:** Mostly found in the thatch layer, chinch bugs infest the base of grass plants. In order to optimize the penetration of the insecticide to location of the chinch bugs, irrigation of the grass prior to treatment may be necessary. If grass is being kept at a long mowing height or if the thatch layer is excessive, use higher volume treatments. It may be necessary to use higher application rates (up to 1 fluid oz. per 1000 square feet) to control populations made up of both adults and nymphs in mid-summer.
- <sup>6</sup> **Mites:** Apply Quali-Pro Bifenthrin I/T 7.9 F in combination with a labeled rate of a surfactant to achieve optimal control of eriophyid mites. A second application may be needed 5 to 7 days after the first to ensure optimal control.
- <sup>7</sup> **Flea Larvae:** Immature fleas mature in shaded areas accessible to pets or other animals. When treating these areas, use a higher volume treatment so that the insecticide penetrates into the soil. NOTE: If adult fleas on lawn areas are being controlled by applying Quali-Pro Bifenthrin I/T 7.9 F at a rate of 0.25 fl. oz. per 1000 square feet, then the rate of larval application can be accomplished by two- to four-fold increase in spray volume.
- <sup>8</sup> **Imported Fire Ants:** The best control will be achieved by using broadcast treatments in combination with mound drenches. This will control present colonies along with foraging workers and newly mated fly-in queens. It is critical either to use high-volume treatments or to irrigate prior to application if the soil is dry. Apply 1 fl. oz. per 1000 square feet when using broadcast treatments. For mound drenches, dilute 1 teaspoon of Quali-Pro Bifenthrin I/T 7.9 F per gallon of water and use 1 to 2 gallons of finished dilution using sufficient force to penetrate the top and allow dilution to flood ant channels. Treat a four-foot diameter around each ant mound. Application should be made in late evening or early morning when it is cooler (65° - 80° when insects are most active). NOTE: A spray rig calibrated to apply 1 fluid oz. per 1000 square feet of Quali-Pro Bifenthrin I/T 7.9 F in 5 gallons per 1000 square feet contains the equivalent dilution (1 teaspoon per gallon) required for fire ant mound drenches in the spray tank.

- <sup>9</sup> **Mole Cricket adults:** Since the preferred grass areas are subject to constant invasion in early spring by the active adult stage, it can be difficult to maintain control of adult mole crickets. It is ideal to treat the areas as late in the day as possible and water immediately after application with up to 0.5 inches of water. To ensure maximum contact when soil is dry, it is necessary to irrigate prior to treatment to bring the adult mole crickets closer to the soil surface. To obtain optimal control of potential nymphal populations, the grass areas preferred by adult mole crickets should be treated at immediately prior to peak hatch stage. (See note 10 below).
- <sup>10</sup> **Mole Cricket nymphs:** Treat grass areas that are preferred by adult mole crickets in the spring just before peak egg hatch. Young nymphs are more vulnerable to insecticidal treatment at this stage because they are close to the soil surface where the insecticide is most concentrated and thereby providing the most efficient control. For larger more damaging nymphal stages later in the year, it may be necessary to use higher application rates more frequently. It is ideal to treat the areas as late in the day as possible and water immediately after application with up to 0.5 inches of water. To ensure maximum contact when soil is dry, it is necessary to irrigate prior to treatment to bring the adult mole crickets closer to the soil surface.
- <sup>11</sup> **Ticks (including ticks that may transmit Lyme Disease and Rocky Mountain Spotted Fever):** Make application to the entire area where contact with ticks may occur. Do not make spot treatments. When applying to areas with heavy leaf litter or dense ground cover, use higher spray volumes. To attain and/or sustain control in times of high pest pressure, re-treatments may be necessary; re-treat only if signs of continued or renewed tick activity are present. Repeat treatments must not be made more often than once per 7 days. **Deer ticks (*Ixodes sp.*)** have a four-stage life cycle spanning 2 years. Treat in late fall and/or early spring to both larval and nymphal stages present in leaf litter and the soil and adults living in the grass and low-lying vegetation above ground. **American dog ticks** invade suburban settings in areas where residences and dwellings are constructed on former fields or wooded areas. These pests normally gather by paths or roadways where they are likely to find a host. To control tick larvae, nymphs and adults, treatments should take place, as needed, from mid-spring to early fall.

### Quali-Pro Bifenthrin I/T 7.9 F Lawn Dilution Chart

<b>Application Volume:</b>	<b>Application Rate:</b>	<b>Fluid Ounces* of Quali-Pro Bifenthrin I/T 7.9 F Diluted to these Volumes of Finished Spray</b>			
<b>Gallons Per 1000 Sq. Ft.</b>	<b>Fluid Ounces per 1000 Sq. Ft.</b>	<b>1 gallon</b>	<b>5 gallons</b>	<b>10 gallons</b>	<b>100 gallons</b>
1.0	0.18	0.18	0.90	1.8	18.0
1.0	0.25	0.25	1.25	2.5	25.0
1.0	0.5	0.5	2.5	5.0	50.0
1.0	1.0	1.0	5.0	10.0	100.0
2.0	0.18	-	0.45	0.90	9.0
2.0	0.25	0.13	0.63	1.25	12.5
2.0	0.5	0.25	1.25	2.5	25.0
2.0	1.0	0.5	2.5	5.0	50.0
3.0	0.18	-	0.30	0.60	6.0
3.0	0.25	-	0.42	0.83	8.3
3.0	0.5	0.17	0.83	1.67	16.7
3.0	1.0	0.33	1.67	3.33	33.3
4.0	0.18	-	0.23	0.45	4.5
4.0	0.25	-	0.31	0.63	6.3
4.0	0.5	0.13	0.63	1.25	12.5
4.0	1.0	0.25	1.25	2.5	25.0
5.0	0.18	-	0.18	0.36	3.6
5.0	0.25	-	0.25	0.5	5.0
5.0	0.5	0.1	0.5	1.0	10.0
5.0	1.0	0.2	1.0	2.0	20.0
10.0	0.18	-	-	0.18	1.8
10.0	0.25	-	0.13	0.25	2.5
10.0	0.5	-	0.25	0.5	5.0
10.0	1.0	0.1	0.5	1.0	10.0

\*To convert to millimeters, multiply by 29.57

1 fluid ounce = 29.57 mL = 2 tablespoons = 6 teaspoons

Do not use household utensils to measure Quali-Pro Bifenthrin I/T 7.9 F.

## ORNAMENTALS AND TREES

Treat with 0.125 to 1.0 fl. oz. of Quali-Pro Bifenthrin I/T 7.9 F per 1000 square feet or 5.4 to 43.5 fl. oz. per 100 gallons for ornamental applications. As long as the highest label rate (1.0 fl. oz. per 1000 square feet or 43.5 fl. oz. per 100 gallons) is not exceeded, Quali-Pro Bifenthrin I/T 7.9 F can be diluted and used in different volumes of water. If diluted with water or other carriers, low-volume equipment can be used for application as long as the highest label rate (1.0 fl. oz. per 1000 square feet or 43.5 fl. oz. per 100 gallons) is not exceeded.

Treat as a full-coverage foliar spray using the stated application rate. If pest pressure and density of foliage increases, repeat treatments using higher rates may be needed to reach the desired control. Repeat treatments should not be made more often than once per 7 days.

Before application to entire planting, test treat a small number of plants and watch for signs of sensitivity. Some plant species may be sensitive to the final spray solution.

To avoid or delay pest resistance, it is recommended to use an alternate class of pesticide in any application program.

## GREENHOUSES AND INTERIOR PLANTSCAPES

Quali-Pro Bifenthrin I/T 7.9 F is effective for use on a broad range of insects and mites on trees, shrubs, leafy plants, nonbearing fruit and nut trees, and flowers grown in greenhouses. It may also be used in interior plants such as hotels, shopping malls, office buildings and other similar areas.

Quali-Pro Bifenthrin I/T 7.9 F may be used by itself or in a tank mix with other products such as insect growth regulators.

### **Calculating Dilution Rates using the Ornamental, Greenhouse, and Interior Plantscape Application Rates Table and the Quali-Pro Bifenthrin I/T 7.9 F Ornamental, Greenhouse, and Interior Plantscape Dilution Chart:**

To determine the proper dilution of Quali-Pro Bifenthrin I/T 7.9 F that is required to control specific pests, follow the steps below:

1. Determine the target pest that is the least susceptible (i.e., the pest that requires the highest application rate for effective control).
2. Choose a treatment rate in terms of fl. oz. of Quali-Pro Bifenthrin I/T 7.9 F.
3. Determine the dilution volume necessary for the treatment.
4. Use the proper amount of Quali-Pro Bifenthrin I/T 7.9 F that must be mixed in your preferred volume of water as shown in the Ornamental Dilution Chart.

As an example, if you were treating for cutworms, the Ornamental Application Rates table shows that 0.125 – 0.25 fluid ounces of Quali-Pro Bifenthrin I/T 7.9 F should be applied per 1000 square feet. Select the application rate of 0.25 fluid oz. per 1000 square feet due to evidence of high pest pressure. The application volume is determined to be 300 gallons per acre which is equivalent to 6.9 gallons per 1000 square feet. The corresponding value in the Ornamental Dilution Chart shows that 0.36 fluid oz. of Quali-Pro Bifenthrin I/T 7.9 F should be mixed with 10 gallons of water.

### **Ornamental, Greenhouse, and Interior Plantscape Application Rates**

Under typical conditions, the application rates in the table below will offer optimal control of the listed pests, but Quali-Pro Bifenthrin I/T 7.9 F can be used at up to 1 fl. oz. per 1000 square feet (43.5 fl. oz. per 100 gallons) at the discretion of the applicator. When maximum residual control is preferred, higher treatment rates are necessary.

**Quali-Pro Bifenthrin I/T 7.9 F Ornamental, Greenhouse, and Interior Plantscape Application Rate Table**

Pest	Application Rate Quali-Pro Bifenthrin I/T 7.9 F	
	Fluid Ounces per 1000 square ft.	Fluid Ounces per 100 gallons
Bagworms <sup>1</sup> Cutworms Elm Leaf Beetles Fall Webworms Gypsy Moth Caterpillars Lace Bugs Leaf Feeding Caterpillars Tent Caterpillars	0.125 - 0.25	5.4 - 10.8
Adelgids <sup>†</sup> Ants Aphids Bees Beet Armyworm Beetles <sup>2,†</sup> Black Vine Weevil (Adults) Brown Soft Scales Broad Mites Budworms California Red Scale (Crawlers) <sup>13</sup> Centipedes Cicadas <sup>†</sup> Citrus Thrips Clover Mites Crickets Diaprepes (Adults) Earwigs European Red Mite Flea Beetles Fungus Gnats (Adults) Grasshoppers Japanese Beetle (Adult) <sup>†</sup> Leafhoppers Leafrollers	0.25 - 0.5	10.8 - 21.7

*(continued)*

**Quali-Pro Bifenthrin I/T 7.9 F Ornamental, Greenhouse, and Interior Plantscape Application Rate Table** (continued)

Pest	Application Rate Quali-Pro Bifenthrin I/T 7.9 F	
	Fluid Ounces per 1000 square ft.	Fluid Ounces per 100 gallons
Mealybugs Millipedes Mites Mosquitoes Orchid Weevil Pillbugs Pine Needle Scales (Crawlers) <sup>2</sup> Plant Bugs (Including <i>Lygus</i> spp.) Psyllids <sup>†</sup> San Jose Scales (Crawlers) <sup>2</sup> Scorpions Sowbugs Spider Mites <sup>3</sup> Spiders Spittlebugs <sup>†</sup> Thrips Tip Moths Treehoppers <sup>†</sup> Twig Borers <sup>2</sup> Wasps Weevils <sup>2</sup> Whiteflies	0.25 - 0.5	10.8 - 21.7
Imported Fire Ants** Leafminers Pecan Leaf Scorch Mite Pine Shoot Beetle (Adults) Spider Mites <sup>14</sup>	0.5 - 1.0	21.7 - 43.5

<sup>1</sup> **Bagworms:** For optimum control, treat when larvae have started to hatch and are young, directing spray to contact as many larvae as possible.

<sup>2</sup> **Beetles, Scale Crawlers, Twig Borers, and Weevils:** Apply to plant foliage; also treat trunks, stems, and twigs.

<sup>3</sup> **Spider Mites:** Apply during spring and mid-summer for most effective control of two-spotted spider mites. During mid- to late-summer, it may be necessary to make more frequent treatments, possibly at higher rates for suitable control. Control may be enhanced by adding a surfactant or horticultural oil or by combining Quali-Pro Bifenthrin I/T 7.9 F with other products registered to control mites. Applications of Quali-Pro Bifenthrin I/T 7.9 F may be alternated with chemicals offering other modes of action that delay or prevent control resistance by two-spotted spider mites. For recommendations on resistance management in your region, check with your local Cooperative Extension Service.

\*\*For foraging ants.

† Not for use in California.

### RESTRICTIONS FOR CALIFORNIA

- Greenhouse applicators must wear: Full-body, chemical-resistant protective suit (such as barrier laminate, butyl rubber, polyvinyl chloride, or equivalent).
- Reapplication Interval: Reapplications to greenhouses must be at intervals of 30 days or longer.
- Greenhouse Harvesters must wear: Regular-length gloves plus a long-sleeved shirt or elbow-length (gauntlet-type) gloves during the 30 days following application.

### Quali-Pro Bifenthrin I/T 7.9 F Ornamental, Greenhouse, and Interior Plantscape Dilution Chart

Application Volume: Gallons Per		Application Rate: Fl. Oz. per	Fluid Ounces* of Quali-Pro Bifenthrin I/T 7.9 F Diluted to these Volumes of Finished Spray			
1000 sq. ft.	Acre	1000 sq. ft.	1 gallon	5 gallons	10 gallons	100 gallons
2.3	100	0.125	-	0.27	0.54	5.4
2.3	100	0.25	0.11	0.54	1.08	10.8
2.3	100	0.5	0.22	1.09	2.17	21.7
2.3	100	1.0	0.44	2.17	4.35	43.5
4.6	200	0.125	-	0.14	0.27	2.7
4.6	200	0.25	-	0.27	0.54	5.4
4.6	200	0.5	0.11	0.54	1.09	10.9
4.6	200	1.0	0.22	1.09	2.17	21.7
6.9	300	0.125	-	-	0.18	1.8
6.9	300	0.25	-	0.18	0.36	3.6
6.9	300	0.5	-	0.36	0.72	7.2
6.9	300	1.0	0.15	0.72	1.45	14.5

\*To convert to millimeters, multiply by 29.57

300 gallons per acre is a typical application volume for landscape ornamental applications.

1 fluid ounce = 29.57 mL = 2 tablespoons = 6 teaspoons

Do not use household utensils to measure Quali-Pro Bifenthrin I/T 7.9 F.

## Pest Control on Outside Surfaces and Around Buildings

Quali-Pro Bifenthrin I/T 7.9 F may be used to control Ants including Carpenter Ants and Fire Ants, Armyworms, Bees, Beetles<sup>†</sup>, Biting Flies, Boxelder Bugs, Centipedes, Chiggers, Chinch Bugs, Clover Mites, Crickets, Cutworms, Dichondra Flea Beetles, Earwigs, Elm Leaf Beetles, Firebrats, Fleas, Flies, Gnats, Grasshoppers, Hornets, Japanese Beetles<sup>†</sup>, Midges, Millipedes, Mosquitoes, Moths, Roaches (including Cockroaches), Scorpions, Silverfish, Sod Webworms, Sowbugs (Pillbugs), Spider Mites, Spiders (including Black Widow Spiders), Springtails, Ticks (including Brown Dog Ticks), and Wasps.

<sup>†</sup> Not for use in California.

Use a 0.02 to 0.06% dilution to spray the outside surfaces of buildings such as private homes, duplexes, townhouses, condominiums, house trailers, apartment complexes, carports, garages, fence lines, storage sheds, barns, and other residential and non-commercial structures. Sites of treatment include but are not limited to exterior siding, foundations, porches, window frames, eaves, patios, garages, garbage areas, lawn areas, trunks of trees and shrubs and other areas where pests may be found. Use a spray volume of up to 10 gallons of emulsion per 1000 square feet. Use higher dilution volumes if vegetation or landscape materials are dense.

Applications to vertical exterior surfaces (e.g., foundations) are permitted to a maximum height of 3 feet from ground level. Sections of vertical exterior surfaces that abut non-porous horizontal surfaces can only be treated if either 1) these sections are protected from rainfall and spray from sprinklers or 2) they do not drain into a sewer, storm drain, or curbside gutter. (e.g., not to sections that abut driveways or sidewalks that drain into streets).

Follow "Additional Application Restrictions for Residential Outdoor Surface and Space Sprays" under **DIRECTIONS FOR USE**.

### Mixing Directions

Suspension	Quali-Pro Bifenthrin I/T 7.9 F per gallon of water	Remarks
0.02%	0.33 fl. oz.	<ul style="list-style-type: none"><li>• Do not use household utensils to measure Quali-Pro Bifenthrin I/T 7.9 F.</li><li>• Use higher treatment rates for quicker knockdown or longer residual control.</li><li>• High pest pressure may require subsequent applications.</li><li>• Repeat application only if there is evidence of renewed insect activity and not more than once per 7 days.</li></ul>
0.06%	1.0 fl. oz.	

**Perimeter Treatment:** Treat a band of soil and vegetation 6 to 10 feet wide around and next to the structure and the foundation of the structure to a height of 2 to 3 feet. Use 0.33 to 1.0 fluid oz. of Quali-Pro Bifenthrin I/T 7.9 F per 1000 square feet in enough water to provide sufficient coverage (refer to Perimeter Application Dilution Chart).

For sections of foundation that abut non-porous horizontal surfaces, the treated areas must be protected from rainfall and spray from sprinklers or they do not drain into a sewer, storm drain, or curbside gutter (e.g., not to sections that abut driveways or sidewalks that drain into streets).

### Quali-Pro Bifenthrin I/T 7.9 F Perimeter Application Dilution Chart

Application Volume: Gallons Per	Application Rate: Fluid Ounces per	Fluid Ounces* of Quali-Pro Bifenthrin I/T 7.9 F Diluted to these Volumes of Finished Spray			
		1000 Sq. Ft.	1 gallon	5 gallons	10 gallons
1	0.33	0.33	1.67	3.33	33.3
1	0.5	0.5	2.5	5.0	50.0
1	0.67	0.67	3.33	6.67	66.7
1	0.75	0.75	3.75	7.5	75.0
1	1.0	1.0	5.0	10.0	100.0
2	0.33	0.17	0.83	1.65	16.5
2	0.5	0.25	1.25	2.5	25.0
2	0.67	0.33	1.67	3.35	33.5
2	0.75	0.38	1.88	3.75	37.5
2	1.0	0.5	2.5	5.0	50.0
3	0.33	0.11	0.55	1.10	11.0
3	0.5	0.17	0.83	1.67	16.7
3	0.67	0.22	1.11	2.23	22.3
3	0.75	0.25	1.25	2.5	25.0
3	1.0	0.33	1.67	3.33	33.3
4	0.33	-	0.41	0.83	8.3
4	0.5	0.13	0.63	1.25	12.5
4	0.67	0.17	0.84	1.67	16.7
4	0.75	0.19	0.94	1.88	18.8
4	1.0	0.25	1.25	2.5	25.0
5	0.33	-	0.33	0.67	6.7
5	0.5	0.1	0.5	1.0	10.0
5	0.67	0.13	0.67	1.33	13.3
5	0.75	0.15	0.75	1.5	15.0
5	1.0	0.2	1.0	2.0	20.0
10	0.33	-	0.17	0.33	3.3
10	0.5	-	0.25	0.5	5.0
10	0.67	-	0.33	0.67	6.7
10	0.75	-	0.38	0.75	7.5
10	1.0	0.1	0.5	1.0	10.0

\*To convert to milliliters, multiply by 29.57

1 fluid oz. = 29.57 mL = 2 tablespoons = 6 teaspoons

**Do not use household utensils to measure Quali-Pro Bifenthrin I/T 7.9 F.**

**Ant and Fire Ant Mound Drench Using 0.06% Dilution:** Use 1-2 gallons of dilution for each mound area. Sprinkle the mound until wet and apply to a 4 ft. diameter circle around the mound. For mounds larger than 12 inches, use a larger volume. Application should be made in late evening or early morning when it is cooler (65° - 80°) when insects are most active.

**Mosquito Control:** To control mosquitoes around buildings, landscapes, and lawns, dilute 0.33 to 1.0 fl. oz. per gallon of water and use one gallon of dilution per 1000 square feet as a general spray. Quali-Pro Bifenthrin I/T 7.9 F can be mixed at lower concentrations and applied at higher volumes to ensure application of the proper amount of product to a given area (refer to the Ornamental or Perimeter Application Charts).

### Other Pest Control Applications

#### Controlling Ants Indoors and Outdoors

**Pest Ants Indoors:** Apply to ant nests for best results. Apply a dilution of 0.5 to 1.0 fl. oz. of Quali-Pro Bifenthrin I/T 7.9 F per gallon of water at the rate of one gallon of dilution per 1000 square feet to places where ants have been seen or are believed to forage as a general surface, spot or crack and crevice treatment. Some of these areas include baseboards, cracks and crevices, in and behind cabinets, under and behind dishwashers, furnaces, refrigerators, sinks and stoves, around pipes, and in corners. Pay close attention when treating entry points into the home or around windows and doors. When combining liquid Quali-Pro Bifenthrin I/T 7.9 F treatments with bait treatments, use Quali-Pro Bifenthrin I/T 7.9 F as instructed above and apply baits in those areas where Quali-Pro Bifenthrin I/T 7.9 F has not been applied.

**Pest Ants Outdoors:** Apply to ant nests for best results. Treat ant trails, around windows and doors, and other places where ants have been seen or are likely to forage. As stated in the "Pest Control on Outside Surfaces and Around Buildings" section, treat using a low- or high-volume perimeter treatment depending on density of vegetation and landscaping materials. When treating concrete surfaces, more frequent treatments, higher dilutions and/or application volumes may be needed for ant control. The following procedures must be followed to help achieve maximum control of the pest:

1. Dilute 0.5 to 1.0 fl. oz. of Quali-Pro Bifenthrin I/T 7.9 F per gallon of water and apply at a rate of up to 10 gallons of dilution per 1000 square feet for maximum residual control.
2. Vegetation and porous surfaces should be treated with high-volume applications using dilutions that are calculated to deliver 0.5 to 1.0 fluid oz. of Quali-Pro Bifenthrin I/T 7.9 F per 1000 square feet (refer to the Ornamental and Perimeter Application Dilution Charts).
3. Treat non-porous surfaces only in areas protected from rainfall and spray from sprinklers with low-volume applications using 0.5 to 1.0 fluid oz. of Quali-Pro Bifenthrin I/T 7.9 F per gallon of water and applying this dilution at the rate of one gallon per 1000 square feet.

**Carpenter Ants Indoors:** Treat areas where carpenter ants are seen or are predicted to forage such as baseboards, in and behind cabinets, under and behind dishwashers, furnaces, refrigerators, sinks, and stoves, around pipes, cracks and crevices, and in corners by diluting 0.5 to 1.0 fluid oz. of Quali-Pro Bifenthrin I/T 7.9 F per gallon of water and applying at the rate of one gallon of dilution per 1000 square feet as a general surface, crack and crevice, spot and/or foam application. Spray or foam into cracks and crevices, or drill holes and spray, mist, or foam into voids and galleries where carpenter ants or their nests are present. When combining liquid Quali-Pro Bifenthrin I/T 7.9 F treatments with bait treatments, use Quali-Pro Bifenthrin I/T 7.9 F as instructed above and apply baits in those areas where Quali-Pro Bifenthrin I/T 7.9 F has not been applied.

**Carpenter Ants Outdoors:** Treat carpenter ant nests for best results. Treat areas where carpenter ants are seen or are believed to forage such as ant trails and around doors and windows. As stated in the “Pest Control on Outside Surfaces and Around Buildings” section, treat using a low- or high-volume perimeter treatment. When treating concrete surfaces, more frequent treatments, higher dilutions and/or application volumes may be needed for carpenter ant control. The following procedures must be followed to help achieve maximum control of the pest:

1. Dilute 0.5 to 1.0 fl. oz. of Quali-Pro Bifenthrin I/T 7.9 F per gallon of water and apply at a rate of up to 10 gallons of dilution per 1000 square feet to obtain residual control.
2. Vegetation and porous surfaces should be treated with high-volume applications using dilutions that are calculated to deliver 0.5 to 1.0 fluid oz. of Quali-Pro Bifenthrin I/T 7.9 F per 1000 square feet (refer to the Ornamental and Perimeter Application Dilution Charts).
3. Treat non-porous surfaces only in areas protected from rainfall and spray from sprinklers with low-volume applications using 0.5 to 1.0 fluid oz. of Quali-Pro Bifenthrin I/T 7.9 F per gallon of water and applying this dilution at the rate of one gallon per 1000 square feet.
4. Use 0.5 to 1.0 fl. oz. of Quali-Pro Bifenthrin I/T 7.9 F per gallon of water on tree trunks with carpenter ant trails or evidence of foraging. Apply to the bark, completely wetting it from the bottom of the tree to the highest possible point on the trunk.

To control carpenter ants inside deck materials, fencing, trees, utility poles or other structural elements, drill to find the inside infested cavity and inject or foam a 0.06% dilution (1.0 fl. oz. of Quali-Pro Bifenthrin I/T 7.9 F per gallon of water) into the cavity with adequate volume and a proper treatment tool with a splash-back guard. Where there are ants tunneling below the surfaces, dilute 0.5 to 1.0 fl. oz. of Quali-Pro Bifenthrin I/T 7.9 F per gallon of water and apply as a drench or foam at intervals of 8 to 12 inches. A uniform barrier should be established where there are ants tunneling below surfaces such as at the edges of walls, driveways or other hard surfaces.

Use a sprinkling can or a hose-end sprayer to distribute a coarse drenching spray; apply a 0.06% dilution to stored lumber and wood piles. This wood may be used for lumber or may be burned after 30 days. Do not use this method of application in structures.

Diluting 1.0 fluid oz. of Quali-Pro Bifenthrin I/T 7.9 F per gallon of water and applying to the soil below where the firewood will be stacked at the rate of one gallon of dilution per 8 square feet will protect the wood from carpenter ants. DO NOT treat firewood with this product.

### **Controlling Termites (Above Ground Only)**

The treatment methods that are expressed below are intended to kill termite workers or winged reproductives present at the time of application. These methods should supplement, not substitute for, mechanical alteration, soil treatment, or foundation treatment.

Controlling winged reproductive termites and exposed workers in localized areas may be accomplished by diluting 1.0 fl. oz. of Quali-Pro Bifenthrin I/T 7.9 F per gallon of water and applying the dilution at the rate of one gallon per 1000 square feet to crawl spaces, unfinished basements, attics, and other crawl spaces as a course fan spray. Both swarming termites and the areas where they gather should be treated.

Controlling above-ground termites in localized areas of infested wood may be accomplished by diluting 1.0 fl. oz. of Quali-Pro Bifenthrin I/T 7.9 F per gallon of water and applying as a foam or a liquid to voids and galleries in wood that is damaged in addition to spaces between wooden structural members and between the foundation and sill plate where the wood is at risk of attack. Drilling and then injecting the foam or dilution into damaged wood or wall voids with an appropriate directional injector will help reach those areas that are not easy to access. After treatment is completed, securely plug the holes that are in regularly occupied areas in the construction elements.

Controlling termite carton nests in building voids can be accomplished by diluting 1.0 fl. oz. of Quali-Pro Bifenthrin I/T 7.9 F per gallon of water and applying as a foam or a liquid using a pointed projection tool. To obtain control, various depths of injection and numerous injection points may be needed. After treatment is complete and when feasible, remove the carton nest material from the building void.

### **Pests Under Slabs**

To control infestations of Arthropods (e.g., ants, cockroaches, and scorpions) that live beneath the slab area, drill or horizontally rod and inject 1 gallon of a 0.06% to 0.12% dilution per 10 square feet or 2 gallons of dilution per 10 linear feet.

### **Posts, Poles, and Other Constructions**

Around wooden constructions (signs, fences, and landscape ornamentation), an insecticidal barrier can be established by treating with a 0.06% dilution. Sub-surface injection and gravity-flow through holes in the bottom of the trench are two treatment methods that can be used on poles and posts that have already been installed. Establishing a complete chemical zone around the pole can be accomplished by treating on all sides. For poles and posts that are fewer than 6 inches in diameter, use 1 gallon of dilution per foot of depth and 1.5 gallons for larger poles applying under the wood to a depth of 6 inches. Four gallons per 10 linear feet per foot of depth should be used for larger constructions.

### **Control of Wood-Infesting Insects in Wood (Localized Areas in Structures)**

<b>Insects</b>	<b>Application Rate</b>	<b>Remarks</b>
Termites Ants Carpenter Ants Wood-infesting beetles (including but not limited to Old House Borer & Powder Post)	Apply a 0.06% dilution to voids and galleries in damaged wood and in spaces between wooden members of a structure and between wood and foundations where wood is at risk.	<ul style="list-style-type: none"> <li>• Can be applied as a paint or fan spray.</li> <li>• Place plastic sheeting under overhead areas that are spot treated except for soil surfaces in crawl spaces.</li> <li>• Areas to which access is difficult can be treated by drilling and then injecting dilution with a crack and crevice injector into the damaged wood or void spaces. (Not intended as a replacement for soil treatment, mechanical alteration, or fumigation to control widespread infestation of wood-infesting insects.)</li> </ul>

### **Control of Wood-Infesting Insects and Nuisance Pests** (Outside of Structures)

In order to control wood-infesting insects active inside trees, utility poles and/or fences, a 0.06% dilution should be injected into the infested cavity which can be found by drilling into the wood. If treating nuisance pests on the exterior of the structure, use a fan spray at a maximum pressure of 25 psi and apply up to the point of runoff. To control Bees, Wasps, Hornets, and Yellow-Jackets, direct the spray at nest openings in the ground, bushes, and in cracks and crevices where the insects may nest. Saturate the openings and contact as many insects as possible.

**Underground Services** (e.g., cables, conduits, pipes, utility lines, wires, etc.) may be in rights-of-way inside of structures or to guard long range (miles) of installations of services.

Treat the soil using a 0.06 to 0.12% Quali-Pro Bifenthrin I/T 7.9 F dilution to prevent and control termite and ant infestations.

Treat the bottom of the trench with 2 gallons of dilution per 10 linear feet and let it soak into the soil. Place the services on the treated soil and cover with about 2 inches of fill soil. Apply another 2 gallons per 10 linear feet over the fill soil to complete the chemical barrier. Only treat the soil in the area near the services in wide trenches, but ensure a continuous barrier of treated soil surrounding the services.

In the event that the soil will not accept the volume stated above, 1 gallon of 0.12% Quali-Pro Bifenthrin I/T 7.9 F may be applied per 10 linear feet of trench over the soil that covers the services and to the base of the trench.

Fill the remainder of the trench with the treated fill soil. Where each service sticks out of the ground, the soil may be treated by trenching/rodding no more than 1 to 2 gallons of dilution into the soil.

**Precautions:** Do not treat electrically active underground services.

## **STORAGE AND DISPOSAL**

Do not contaminate water, food or feed by storage or disposal.

**STORAGE AND SPILL PROCEDURES:** Store upright at room temperature. Avoid exposure to extreme temperatures. In case of spillage or leakages, soak up with an absorbent material such as sand, sawdust, earth, Fuller's earth, etc. Dispose of with chemical waste.

**PESTICIDE DISPOSAL:** Pesticide, spray mixture, or rinse water that cannot be used according to label instructions must be disposed of at or by an approved waste disposal facility.

### **CONTAINER DISPOSAL:**

**For Containers equal to or less than 5 Gallons:** Nonrefillable container. Do not reuse or refill this container. Clean container promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container ¼ full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Offer for recycling if available. If recycling is not available, puncture or dispose of in a sanitary landfill or incineration or if allowed by state and local authorities, by burning. If burned stay out of smoke.

## LIMITATION OF WARRANTY AND LIABILITY

Read the entire directions for use, conditions of warranties and limitations of liability before using this product. If terms are not acceptable, return the unopened product container at once.

By using this product, user or buyer accepts the following **Conditions, Disclaimer of Warranties and Limitations of Liability**.

**CONDITIONS:** The directions for use of this product are believed to be adequate and must be followed carefully. However, it is impossible to eliminate all risks associated with the use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or the manner of use or application, all of which are beyond the control of Control Solutions, Inc. All such risks shall be assumed by the user or buyer.

**DISCLAIMER OF WARRANTIES:** To the extent consistent with applicable law, Control Solutions, Inc. makes no other warranties, express or implied, of merchantability or of fitness for a particular purpose or otherwise, that extend beyond the statements made on this label. No agent of Control Solutions, Inc. is authorized to make any warranties beyond those contained herein or to modify the warranties contained herein. To the extent consistent with applicable law, Control Solutions, Inc. disclaims any liability whatsoever for special, incidental or consequential damages resulting from the use or handling of this product

**LIMITATIONS OF LIABILITY:** To the extent consistent with applicable law, the exclusive remedy of the user or buyer for any and all losses, injuries or damages resulting from the use or handling of this product, whether in contract, warranty, tort, negligence, strict liability or otherwise, shall not exceed the purchase price paid or at Control Solutions, Inc.'s election, the replacement of product.

Micro-injector is a registered trademark of Whitmire Micro-Gen Research Laboratories.

Actisol is a registered trademark of Roussel-Uclaf.

Quali-Pro is a registered trademark of Makhteshim Agan of North America, Inc.





# QUALI-PRO<sup>®</sup>

## Bifenthrin I/T 7.9 F

### *Insecticide/Termiticide*

For prevention and control of listed termites carpenter ants and other pests of structures. To control listed pests in and around such areas as homes, commercial and industrial buildings, recreational areas, athletic fields, lawns, ornamentals, and greenhouses. Controls pests in livestock and poultry houses.

For Use as a Termiticide: May only be used by individuals/firms licensed by the State to apply termiticide products. States may have more restrictive requirements regarding qualifications of persons using this product. Consult the pest control regulatory agency of your State prior to use of this product.

ACTIVE INGREDIENT:	% BY WT.
Bifenthrin* .....	7.9%
INERT INGREDIENTS: .....	92.1%
TOTAL: .....	100.0%

Quali-Pro Bifenthrin I/T 7.9 F contains 2/3 pound active ingredient per gallon.

\*Cis isomers 97% minimum, trans isomers 3% maximum.

Contains bifenthrin, the active ingredient used in Talstar P Professional Insecticide.

Quali-Pro Bifenthrin I/T 7.9 F is not manufactured or distributed by FMC Corp.

EPA Reg. No. 53883-365

**KEEP OUT OF REACH OF CHILDREN**  
**CAUTION**

For additional precautionary, handling, and use statements, see inside of this booklet.

Manufactured for:  
**Control Solutions Inc.**

5903 Genoa Red Bluff • Pasadena, Texas 77507



EPA 20150225

**Net Contents: 1 Gallon**

EPA Est. No. 53883-TX-002<sup>CS</sup>; 37429-GA-001<sup>BT</sup>; 37429-GA-002<sup>BO</sup>  
Letter(s) in lot number correspond(s) to superscript in EPA Est. No.